

Fax Number: 780-814-5593

A division of



## **REQUEST FOR PULMONARY FUNCTION TESTING**

Address: #108 10814 100 Street Grande Prairie, Alberta T8V 2M8 Phone:780-814-5563	
Patient Information	Physician Information
Last Name:	Referring Clinic:
First Name:	Referring Physician:
	Practitioner ID:
DOB: // PHN:	Fax: Phone:
Gender Identity:	
☐ Male ☐ Female ☐ Non-Binary	Family Physician:
Transgender Male Transgender Female	Fax: Phone:
Prefer to Describe	Referral Date:
Home Address:	
	History / Reasons for Referral
City	Asthma Cough
City:	☐ Dyspnea ☐ Wheeze
Province:Postal Code:	Fibrosis COPD
Phone Number:	Other:
Procedu	re Request
	Full Pulmonary Function Test
	Spirometry
PFT Referral Guidelines for Physicians	PFT Referral Guidelines and Pre -Test Instructions for Patients
Definitions     Full PFT : Pre & Post Bronchodilator Spirometry / Diffusion Capacity / Lung Volumes	Do not take ( <u>if possible</u> ) the following medications prior to your appointment:
	Inhaled bronchodilators:
Spirometry : Pre & Post Bronchodilator Spirometry	• Short-acting for 4 hours: (e.g. albuterol, salbutamol, ventolin, bricanyl)
Conditions where suboptimal lung function results or test performance are likely:	Long-acting for 24 hours (e.g. formoterol, salmeterol, Oxeze, Serevent, Symbicort, Advair)
Chest or abdominal pain     Oral or fiscal pain by mouthpiece.	Ultra long-acting agents for 36 hours (e.g. Onbrez, Breo, Inspiolto, Anoro)
Oral or facial pain by mouthpiece     Stress incontinence	Long Acting Muscarinic Antagonists 36-48 (e.g. Spiriva, Incruse, Tudorza, Seebri, Anoro,
4. Dementia or confused state	Inspiolto, Ultibro, Duaklir Anticholinergics:
5. Inability to follow instructions (e.g., < 6 years, Mental Health Condition)	Short-acting for 12 hours (e.g. Atrovent)
Other contraindications and waiting period before testing include:  1. Recent eye surgery 1 week to 6 months (depending on type of surgery)	Long-acting for 36-48 hours (e.g. Tiotropium)
Recent eye surgery 1 week to 6 months (depending on type of surgery)     Recent brain surgery or injury (4 weeks)	Take as Usual
3. Pneumothorax (6 weeks)	Take Theophylline preparations and oral steroids as usual
Hemoptysis of unknown origin or related to transmissible respiratory infection     Current Pneumothorax	You should not have a Pulmonary Function Test if you have had:
Current Predmotroiax     Unstable cardiovascular status / Acute MI / Non-Compensated Heart Failure - (1 week)	A) Pneumothorax (Lung Collapse) in the last 6 weeks B) Eye Surgery (including Laser Surgery)
7. Thoracic, abdominal, or cerebral aneurysms	in the last 1 week to 6 months (depending on type of eye surgery)  C) Surgery of the chest, neck or abdomen in the last 4 weeks D) Unstable cardiovascular
8. Active or suspected transmissible respiratory infections 9. Sinus or middle ear surgery within - (1 week)	C) Surgery of the chest, neck of abdomen in the last 4 weeks D) Unstable cardiovascular condition within 1 week. E) Any type of aneurysm

9. Sinus or middle ear surgery within - (1 week)

10. Abdominal or Thoracic surgery (4 weeks)

11. Late term Pregnancy

\*\*Refrain from smoking for 3 hours prior to testing\*\*

\*\*Refrain from taking anything containing caffeine 3 hours prior to testing\*\*